

**THE SOUTH PACIFIC SEA LEVEL & CLIMATE
MONITORING PROJECT**

MONTHLY DATA REPORT

NO. 170

AUGUST 2009



Australian Government

Bureau of Meteorology

This project is sponsored by the Australian Agency for International Development (AusAID), and is managed by the Bureau of Meteorology with its National Tidal Centre (NTC) providing key technical support.



Australian Government

Bureau of Meteorology

National Tidal Centre Bureau of Meteorology Australia

GPO Box 421
Kent Town SA 5071
Australia

Tel: (+618) 8366 2730
Fax: (+618) 8366 2651
Website: <http://www.bom.gov.au/oceanography/>

Quality Certification:

I authorise the issue of this South Pacific Sea Level and Climate Monitoring Project Monthly Data Report for August 2009 in accordance with National Tidal Centre Quality Assurance procedures.

William Mitchell
Manager - National Tidal Centre

South Pacific Sea Level and Climate Monitoring Project

Monthly Data Report

August 2009

EXECUTIVE SUMMARY

This summary, and the overview that follows, are intended to provide a synopsis of the Monthly Data Report and of the trends observed over the life of the project to date.

August 2009

- The SEAFRAME network continued to collect high quality sea level and associated meteorological information for monitoring climate variability and climate change.
- Sea levels during August were generally close to what is normally observed at this time of the year. Disruptions to annual sea level cycles associated with El Niño indicators have not yet been observed, but will be monitored closely in the coming months.
- Warmer than normal ocean temperatures and weaker than normal Trade Winds continued to be observed across the equatorial Pacific. However abnormally warm sea surface temperatures in the western equatorial Pacific and subsurface cooling has hindered the development of atmospheric circulation typical of El Niño.
- The majority of international climate models predict equatorial Pacific surface temperatures will remain warmer than normal for the remainder of 2009. The likelihood of the maturing of El Niño conditions during 2009 remains high.

Short-Term Trends

It is important to stress that as the sea level record becomes longer, the short-term trend estimate becomes more stable and reliable. Observed trends in sea level include natural variability, for example, events such as El Niño and effects due to many other atmospheric, oceanographic and geological processes. Longer-term data sets for all stations are required in order to separate the effects of the different signals. ***Please exercise caution in interpreting the short-term trends in the table below*** – they will almost certainly change over the coming years as the data set increases in length. Figure 13 later in this report provides the “time history” of the short-term trend at all project locations.

Recent short-term sea level trends in the project area based upon SEAFRAME data through August, 2009				
Location	Lat / Long	Installation Date	Trend (mm/yr)	Change from previous month
Cook Is	21°12'17.1"S / 159°47'5.2"W	Feb 1993	+5.3	0.0
Tonga	21°8'12.5"S / 175°10'50.5"W	Jan 1993	+9.7	+0.1
Fiji	17°36'17.7"S / 177°26'17.7"E	Oct 1992	+5.7	+0.1
Vanuatu	17°45'19.2"S / 168°18'27.7"E	Jan 1993	+6.1	+0.1
Samoa	13°49'36.4"S / 171°45'40.7"W	Feb 1993	+5.9	-0.1
Tuvalu	8°30'8.9"S / 179°11'42.6"E	Mar 1993	+5.3	-0.1
Kiribati	1°21'54.2"N / 172°55'58.8"E	Dec 1992	+3.5	+0.2
Nauru	0°31'45.9"S / 166°54'36.2"E	Jul 1993	+4.4	+0.1
Solomon Is.	9°25'44.1"S / 159°57'19.3"E	Jul 1994	+8.5	-0.1
PNG	2°2'31.5"S / 147°22'25.6"E	Sep 1994	+8.0	-0.1
FSM	6°58'49.9"N / 158°12'0.8"E	Dec 2001	+19.5	-0.4
Marshall Is.	7°6'21.7"N / 171°22'22.1"E	May 1993	+4.2	0.0

INTRODUCTION

Welcome to the August 2009 Monthly Data Report for the South Pacific Sea Level and Climate Monitoring Project (SPSLCMP). The report details the month by month operation of the SEAFRAME monitoring stations in the Pacific, including operational problems with the network or with satellite communications, the occurrence of abnormal sea level or climate events, interpretation of sea level fluctuations in the context of El Niño and the emergence of trends in the data.

The SPSLCMP was developed as an Australian response to concerns raised by the member countries of the South Pacific Forum over the potential impacts of global warming on climate and sea levels in the Pacific. Support was provided for the installation of SEAFRAME monitoring stations across the South Pacific Forum region.

SEAFRAME gauges not only measure sea level by two independent means, but also observe a number of “ancillary” variables - air and water temperatures, wind speed, wind direction and atmospheric pressure. There is an associated programme of levelling to first order, to determine shifts in the vertical of the sea level sensors due to local land movement. Continuous Global Positioning System (CGPS) measurements are now also being made to determine the vertical movement of the land with respect to the International Terrestrial Reference Frame.

The AusAID funded project has, as its principal objective *‘the provision of an accurate long term record of sea level in the South Pacific for partner countries and the international scientific community, that enables them to respond to and manage related impacts’*.

The project's monitoring network consists of 12 SEAFRAME stations, providing a wide coverage across the Southwest Pacific basin. All of these stations (see Figure A), with the exception of the Pohnpei (FSM) gauge, which was established in December 2001, have been operational since October 1994.

The monthly data report, one of a range of information products produced by the project, is the primary form of SPSLCMP data dissemination. Its content is designed to provide up-to-date access to the project's data products.




Figure A: *South Pacific Sea Level and Climate Monitoring Stations*

AUGUST CLIMATOLOGY

Climate conditions across the equatorial Pacific during August resembled a weak but unusual El Niño. Ocean heat content across the central and eastern equatorial Pacific remained warmer than normal and continued to exceed El Niño thresholds but the far western tropical Pacific and Coral Sea is also warmer than average, which is unusual for El Niño. As a consequence atmospheric indicators such as the Southern Oscillation Index, Trade Winds and equatorial cloudiness are yet to show patterns typical of an El Niño. The majority of international climate models predict that sea surface temperatures across the equatorial Pacific will remain above El Niño thresholds for the remainder of 2009.

The Southern Oscillation Index (SOI) is usually consistently below -7 during El Niño events but the August value of -5 following the July value of $+2$ is indicative of near neutral conditions (**Figure B**).

Sea surface temperatures across the central and eastern equatorial Pacific were warmer than normal during August and exceeded El Niño thresholds (**Figure C**). Anomalies in excess of $+1^{\circ}\text{C}$ were observed in these areas with little change to the overall sea surface warmth since July.

Subsurface ocean temperatures were also warmer than average across the central and eastern equatorial Pacific Ocean during August, with anomalies exceeding $+2.0^{\circ}\text{C}$ in some areas. However subsurface ocean heat content has been cooling steadily since June (**Figure D**).

During El Niño (warm-episode) conditions there is a sustained weakening of the Trade Winds across much of the equatorial Pacific and an increase in cloudiness in the central Pacific particularly near the dateline. During La Niña (cold-episode) conditions there is a reversal of this situation, with stronger Trade Winds and a decrease in cloudiness in the central Pacific. During August 2009 the easterly Trade Winds were weaker than normal over most of the equatorial Pacific (**Figure E**). Cloudiness over the equatorial Pacific near the dateline has increased in recent months, although for the first 3 weeks of August cloudiness was actually near to below average. The El Niño trend of increasing cloudiness near the dateline has been relatively weak by comparison to other events.

The majority of seven international computer models surveyed by the Bureau of Meteorology predict sea surface temperatures will remain above El Niño thresholds for the remainder of 2009. While the extent of warming is more moderate than in previous forecasts, the likelihood of an El Niño event continuing to develop through 2009 remains high.

The preceding description of the climatology of the Pacific region, and Figures B, C and D are based on information sourced from the National Climate Centre of the Australian Bureau of Meteorology at <http://www.bom.gov.au/climate/>. Figure E was generated from the Tropical Atmosphere Ocean project website courtesy of PMEL, NOAA at <http://www.pmel.noaa.gov/tao/>.




Figure B: The five-month weighted mean and individual monthly means of the Southern Oscillation Index (SOI). The SOI is ten times the monthly anomaly of the difference in mean sea level pressure between Tahiti and Darwin, divided by the long-term standard deviation of that difference for the relevant month.




Figure C: Sea surface temperature anomaly (°C) for August 2009.




Figure D: Equatorial depth-longitude section of ocean temperature anomalies for May 2009 through to August 2009. Contour interval is 0.5°C .




Figure E: Monthly mean wind vectors (top) and anomalies (bottom) for August 2009. The colour-shaded contours represent the monthly mean sea surface temperatures (top) and anomalies (bottom).

AUGUST SEAFRAME DATA

Monthly Sea Level and Environmental Data (Figures 1-10)

The **observed sea levels (Figure 1)** are dominated by the daily oscillations of the tide. In most cases, the tide rises and falls twice per day (semi-diurnal), but at PNG and the Solomon Islands the tide tends to have a single high and low per day (diurnal). The greatest variations are called spring tides and tend to occur close to the full and new moon. There was a full moon on the 6th of August and a new moon on the 20th of August UTC.

Gaps in the data are the result of instrumental errors or data retrieval problems and are discussed under **Instrument Performance**.

The **residuals (Figure 2)** are the differences between the observed sea levels and the tidal predictions. They highlight the non-tidal sea level fluctuations such as those due to the short-term effects of weather or tsunamis. Residual sea level fluctuations may also be amplified or sustained by the shape of the harbour in which the gauge is located. Persistent sloshing of water within a bay or harbour, for example, is known as a seiche. Seiches are often recorded at PNG when the wind suddenly changes strength or direction. Large non-tidal sea level fluctuations are sometimes observed at FSM during periods of reduced or neap tides, such as occurred from 24th – 30th August 2009.

The sea level residuals at all stations, to some degree, exhibit semi-diurnal or diurnal fluctuations, which last a few days or weeks and then disappear. If these peaks were to persist, rather than appear as occasional ‘transients’, then the tidal analysis would be able to account for them, and the end result would be virtual eradication from the residuals.

The **barometrically corrected residuals (Figure 3)** have had the effect of atmospheric pressure fluctuations removed from the sea level residuals of Figure 2. The rule of thumb for the ‘inverse barometer effect’ is that a 1-hPa fall in the barometer, if sustained over a day or more, produces a 1 cm rise in the local sea level (within the area beneath the low pressure system).

The **winds, temperatures and barometric pressures** are plotted in **Figures 4 to 9**. The short lines in **Figure 5** follow the meteorological convention, that is, they point in the direction the wind is coming *from*. For example, the winds at Tuvalu prevailed from the southeast for most of the month.

Air and water temperatures (**Figures 7 and 8**) are plotted using the same vertical scale for the purpose of comparison. The air temperatures are seen to fluctuate over a much wider range than the water temperatures. At some sites (e.g. FSM) the water temperature shows almost no variation, although the air temperature varies by several degrees between night and day. At Nauru a twice-daily fluctuation in water temperature is related to the tide, as it is usually more pronounced during the larger spring tides.

Barometric pressures (**Figure 9**) tend to fluctuate by around 3 hPa twice-daily at all stations as a result of atmospheric tides, which are largest in the tropical regions and reduce to near zero toward the poles. The longer-term barometric pressure fluctuations

that occur over periods of days to weeks are due to passing weather systems. These fluctuations tend to be larger at sites further away from the equator such as Cook Islands and Tonga.

The **meteorological data** are put into perspective by **Figure 10**. In this figure, if an open circle falls above (below) a solid dot, a new maximum (minimum) for the particular month has been set. *The data sets only include South Pacific Sea Level and Climate Monitoring Project data, which have been collected since October 1992 when the first station was installed (Fiji). The data from FSM has only been collected since December 2001.* A new maximum August air temperature of 30.0°C was recorded at PNG.

Mean Sea Level and Anomalies (Figures 11-13)

Figure 11 shows the **monthly mean sea levels**, which are simple arithmetic averages of the sea levels, relative to an arbitrary zero. The figure shows that Tuvalu, for example, normally experiences an annual cycle of about 0.2 metres, reaching a peak around February or March. One effect of the El Niño of 1997/1998 was to disrupt the annual sea level cycle at many of the SEAFRAME stations.

Figure 12 shows the **monthly mean sea level anomalies**, or departures from normal conditions after tides, annual and semi-annual seasonal cycles and the sea level trend have been removed. The annual cycle at Tuvalu (which has the largest consistent annual cycle) is quite notable in **Figure 11** but less apparent in **Figure 12**. By removing the seasonal cycles, the anomalies help to bring out irregular features, such as lower than normal sea levels across the region during the 1997/98 El Niño.

In August 2009 sea levels were 5-10cm higher than normal at Kiribati, Nauru, Vanuatu, Fiji and Tonga. Slightly lower than normal sea levels were observed at FSM, PNG, Solomon Islands, Tuvalu and Samoa. Lower than normal sea levels typically associated with El Niño conditions are yet to materialise in the region.

Sea Level Trends

The **short-term sea level trends** at individual stations as at August 2009 are shown in the following table. Sea level trends are updated every month by allowing for a linear trend term in the tidal analysis of all the data available at individual stations. *Please exercise caution in interpreting the trends* – they will continue to change over the coming years as the data sets increase in length. The evolution of the monthly trend values (in mm per year) at each station from one year after installation to present is depicted in **Figure 13**. This figure illustrates that as the sea level record becomes longer, the relative sea level trend estimates become more stable and reliable. The reason for this is that the trends from short sea level records are affected by the natural sea level variability occurring on inter-annual, El Niño and decadal timescales due to atmospheric, oceanographic and geological processes. Longer-term data sets for all stations are required in order for the underlying trend to emerge from these short-term variations. Further details are available from the *National Tidal Centre (NTC), Australian Bureau of Meteorology*.

Recent short-term sea level trends in the project area based upon SEAFRAME data through August, 2009				
Location	Lat / Long	Installation Date	Trend (mm/yr)	Change from previous month
Cook Is	21°12'17.1"S / 159°47'5.2"W	Feb 1993	+5.3	0.0
Tonga	21°8'12.5"S / 175°10'50.5"W	Jan 1993	+9.7	+0.1
Fiji	17°36'17.7"S / 177°26'17.7"E	Oct 1992	+5.7	+0.1
Vanuatu	17°45'19.2"S / 168°18'27.7"E	Jan 1993	+6.1	+0.1
Samoa	13°49'36.4"S / 171°45'40.7"W	Feb 1993	+5.9	-0.1
Tuvalu	8°30'8.9"S / 179°11'42.6"E	Mar 1993	+5.3	-0.1
Kiribati	1°21'54.2"N / 172°55'58.8"E	Dec 1992	+3.5	+0.2
Nauru	0°31'45.9"S / 166°54'36.2"E	Jul 1993	+4.4	+0.1
Solomon Is.	9°25'44.1"S / 159°57'19.3"E	Jul 1994	+8.5	-0.1
PNG	2°2'31.5"S / 147°22'25.6"E	Sep 1994	+8.0	-0.1
FSM	6°58'49.9"N / 158°12'0.8"E	Dec 2001	+19.5	-0.4
Marshall Is.	7°6'21.7"N / 171°22'22.1"E	May 1993	+4.2	0.0

Barometric Pressure, Water Temperature and Air Temperature Anomalies

The anomalies of barometric pressure, water and air temperature (**Figures 14 to 16**) are determined in the same manner as the sea level anomalies (**Figure 12**), except the trend is not calculated.

The **barometric pressure anomalies** (**Figure 14**) show substantially higher than normal barometric pressures were observed at SEAFRAME stations during the 1997-1998 El Niño. In August 2009 near normal barometric pressures continued to be observed across the region and are yet to show any regional pattern associated with the developing El Niño conditions.

The **water temperature anomalies** (**Figure 15**) show slightly warmer than normal water temperatures were observed in August 2009 at Marshall Islands, FSM, PNG, Solomon Islands, Kiribati and Nauru. Elsewhere sea levels were near to what is normally observed at this time of the year.

The **air temperature anomalies** (**Figure 16**) show conditions during August 2009 were generally near to what is normally observed at this time of the year, with the largest anomaly of +0.7°C associated with warmer than normal conditions at Marshall Islands. Over the duration of the record the air temperature anomalies generally (although not always) follow the water temperature anomalies, which is an indication of the large influence the ocean has upon the climate of the Pacific Islands.

Instrument Performance

In **Figure 17**, which shows **sea level data return**, colour is used to distinguish five-year project phases. The number of missing days is noted in gaps in the bars.

Sea level data return was good at most stations during August 2009 with the exception of PNG, where problems with both satellite and phone communications caused data losses totalling 7 days during the month. At Nauru problems with the primary sea level sensors were encountered and data from the secondary sea level sensors were used where possible.


Various problems were encountered with ancillary meteorological sensors including the air temperature, water temperature and barometric pressure electronic circuit at Marshall Islands and the resulting erroneous data were removed from the record. The water temperature sensor at Tonga experienced problems and likewise erroneous data were removed.

SEAFRAME STATIONS

SEAFRAME stations employ a SUTRON programmable data logger, water level gauges and other sensors. The data logger and associated electronics are normally housed in fibreglass huts. A sketch of a typical station is shown in the following figure. Water level sensors include:


- (1) Primary water level using a Bartex 'AQUATRAK' acoustic-in-air sensor, and
- (2) Secondary water level (or backup) using a Druck pressure transducer mounted close to the seabed.

The primary and backup water level sensors provide water level values, which are averaged over three minutes and are logged every six minutes. The data logger has the memory capacity to store approximately one month of data. The meteorological sensors are logged to the SUTRON data logger on an hourly basis.



Web Hits

The following chart shows the number of times the Pacific pages on the *NTC* web site have been visited, by month since January 2006.



The *Monthly Data Report* is prepared by *NTC* for *AusAID*.

NTC would appreciate feedback from readers on the content and presentation of the *Monthly Data Report*.

Please spare a few moments to let us know your constructive opinion.

Further communication on the *Monthly Data Report* may be made to *NTC*. Anyone interested in a more detailed account of the project should contact:

National Tidal Centre
Bureau of Meteorology
PO Box 421
Kent Town SA 5067
Tel: (+618) (08) 8366 2600
Fax: (+618) (08) 8366 2693
Website: <http://www.bom.gov.au/oceanography/tides.shtml>

Or visit the project website at <http://www.bom.gov.au/pacificsealevel>

Please refer to: <http://www.bom.gov.au/oceanography/projects/spslcmp/spslcmp.shtml> for details.

Please also note the following:

While care has been taken in the collection, analysis, and compilation of the data, it is supplied on the condition that neither the *Commonwealth of Australia* nor *NTC* shall be liable for any loss or injury whatsoever arising from the use of the data. Copyright for material contained in this document is held by the *Commonwealth of Australia*.

Individuals and organisations are advised that quality controlled six-minute or hourly data from these stations are available on request from *NTC*. Some handling fees may be charged. For commercial agencies requesting data, some additional costs may be levied.

Figure 1

AUGUST 2009

SIX MINUTE WATER LEVEL OBSERVATIONS (m)




Figure 2

AUGUST 2009
SIX MINUTE RESIDUAL WATER LEVELS (m)




Figure 3

AUGUST 2009

SIX MINUTE RESIDUALS

ADJUSTED FOR ATMOSPHERIC PRESSURE (m)




Figure 4

AUGUST 2009
HOURLY WIND SPEEDS (m/s)




Figure 5
AUGUST 2009
HOURLY INCIDENT WINDS (m/s, deg True)




Figure 6
AUGUST 2009
HOURLY MAXIMUM WIND GUSTS (m/s)




Figure 7

AUGUST 2009
HOURLY AIR TEMPERATURES (°C)




Figure 8

AUGUST 2009
HOURLY WATER TEMPERATURES (°C)




Figure 9
AUGUST 2009
HOURLY ATMOSPHERIC PRESSURE (hPa)







Figure 10

Comparison of August 2009 Max, Min & Mean with Long Term August Values



Stations

- 1 - Marshall Islands
- 2 - Federated States of Micronesia
- 3 - Papua New Guinea
- 4 - Solomon Islands
- 5 - Kiribati
- 6 - Nauru
- 7 - Tuvalu
- 8 - Samoa
- 9 - Vanuatu
- 10 - Fiji
- 11 - Tonga
- 12 - Cook Islands

- August 2009 Maximum
- August 2009 Mean
- August 2009 Minimum
- Long Term August Maximum
- Long Term August Mean
- Long Term August Minimum

MONTHLY MEAN SEA LEVELS TO AUGUST 2009 (m)

The chart displays the annual percentage change in the population aged 65 and over for 12 Pacific Island countries from 1992 to 2009. The countries are listed vertically: Marshall Islands, Federated States of Micronesia, Papua New Guinea, Solomon Islands, Kiribati, Nauru, Tuvalu, Samoa, Vanuatu, Fiji, Tonga, and Cook Islands. Each country's data is represented by a line with markers. The y-axis for each panel ranges from -0.2 to 0.2, with a horizontal line at 0. The x-axis for all panels shows years from 1992 to 2009. The data shows varying trends, with some countries like Samoa and Tuvalu showing significant fluctuations and others like the Cook Islands showing more stable, lower growth rates.

Figure 12
SEA LEVEL ANOMALIES THROUGH AUGUST 2009 (m)




Figure 13

SEA LEVEL TRENDS THROUGH AUGUST 2009 (mm/year)




Figure 14

BAROMETRIC PRESSURE ANOMALIES THROUGH AUGUST 2009 (hPa)




Figure 15
**WATER TEMPERATURE ANOMALIES
THROUGH AUGUST 2009 (°C)**




Figure 16
**AIR TEMPERATURE ANOMALIES
THROUGH AUGUST 2009 (°C)**




Figure 17

SEA LEVEL DATA RETURN

THE NUMBER OF DAYS OF GAP ARE INDICATED

GAPS INCLUDE TRANSMISSION, POWER AND LOGGER FAILURE

* Patchy record

