



Climate and Oceans Support
Program in the Pacific



Australian Government
Bureau of Meteorology

Monthly Data Report - October 2012

Pacific Sea Level Monitoring Project





Australian Government
Bureau of Meteorology

© Commonwealth of Australia 2012

This work is copyright. Apart from any use as permitted under the *Copyright Act 1968*, no part may be reproduced without prior written permission from the Bureau of Meteorology. Requests and inquiries concerning reproduction and rights should be addressed to the Publishing Unit, Bureau of Meteorology, GPO Box 1289, Melbourne 3001. Requests for reproduction of material from the Bureau website should be addressed to AMDISS, Bureau of Meteorology, at the same address.

Published by Bureau of Meteorology



Table of Contents

List of Tables	iv
List of Figures	iv
Executive Summary.....	1
Introduction.....	2
Sea Level and Climate	4
October SEAFRAME Data	5
Monthly Sea Level and Environmental Data	5
Mean Sea Level and Anomalies	7
Overall Rate of Movement in Sea Level.....	8
Barometric Pressure, Water Temperature and Air Temperature Anomalies.....	9
Instrument Performance	10
SEAFRAME Stations.....	11
Observation Network Upgrade Project.....	12
Further Information	13
Online Resources	13
Acknowledgement	13
Appendix 1: SEAFRAME Data Figures	14

List of Tables

Table 1. Updated overall rates of sea level movement based on SEAFRAME data from installation through October, 2012.....	8
Table 2. Rates of sea level data return.....	10
Table 3. Schedule of SEAFRAME station equipment upgrades.....	12

List of Figures

Figure 1. Pacific Sea Level Monitoring Network of SEAFRAME stations	3
Figure 2. Schematic diagram of a SEAFRAME sea level monitoring station.....	11
Figure 3. Sea level observations during October 2012.....	15
Figure 4. Residual sea levels during October 2012.....	16
Figure 5. Residual sea levels adjusted for atmospheric pressure during October 2012.....	17
Figure 6. Wind speeds during October 2012.....	18
Figure 7. Incident winds during October 2012.....	19
Figure 8. Wind gusts during October 2012.....	20
Figure 9. Air temperatures during October 2012.....	21
Figure 10. Water temperatures during October 2012.....	22
Figure 11. Atmospheric pressures during October 2012.....	23
Figure 12. Comparison of October 2012 ancillary data with long term October values	24
Figure 13. Monthly mean sea levels to October 2012	25
Figure 14. Monthly sea level anomalies to October 2012.....	26
Figure 15. Monthly barometric pressure anomalies to October 2012.....	27
Figure 16. Monthly water temperature anomalies to October 2012.....	28
Figure 17. Monthly air temperature anomalies to October 2012.....	29
Figure 18. Sea level data return.....	30



Executive Summary

This summary, and the overview that follows, are intended to provide a synopsis of the recent month's observations in addition to longer term variations over the life of the project to date.

October 2012

- The SEAFRAME network continued to collect high quality sea level and associated meteorological information for monitoring climate variability and climate change.
- Significant station outages were encountered at FSM and Cook Islands. The station at Cook Islands was upgraded and returned to service with a modern data logger, improved real time communications capability and an additional radar-type sea level sensor during October 2012.
- Monthly sea levels during October were about 5cm higher than normal at Kiribati, Nauru and Fiji and 5cm lower than normal at Solomon Islands, Tuvalu and Vanuatu. At Marshall Islands, PNG, Samoa, Tonga and Cook Islands monthly sea levels were near normal for this time of the year.
- Sea levels remained close to predicted astronomical tides on a daily basis throughout October. No significant storm surge was observed.
- Small tsunami signals on the order of 10cm were observed at Cook Islands and Samoa following a magnitude Mw7.7 earthquake on October 28 in the northeast Pacific off the coast of Canada.

Introduction

Welcome to the October 2012 Monthly Data Report for the Pacific Sea Level Monitoring Project (PSLMP). The report details the month by month operation of the SEAFRAME monitoring stations in the Pacific, including operational problems with the network or with satellite communications, the occurrence of abnormal sea level events and the interpretation of sea level fluctuations in the context of related astronomical tide, weather and climate variations.

The PSLMP continues the work of the South Pacific Sea Level and Climate Monitoring Project (SPSLCMP) under a wider Climate and Oceans Support Program in the Pacific (COSPPac) initiative. The SPSLCMP was originally developed as an Australian response to concerns raised by the member countries of the South Pacific Forum over the potential impacts of global warming on climate and sea levels in the Pacific with the principal objective of 'the provision of an accurate long term record of sea level in the South Pacific for partner countries and the international scientific community which enables them to respond to and manage related impacts'.

The project's sea level monitoring network consists of 12 SEAFRAME stations providing wide coverage across the Pacific Islands Forum region (Figure 1). The SEAFRAME stations not only measure sea level, but also observe a number of "ancillary" variables - air and water temperatures, wind speed, wind direction and atmospheric pressure.

An associated geodetic measurement program, implemented by Geosciences Australia, supports levelling surveys to first order, to determine shifts in the vertical of the sea level sensors due to local land movement, as well as continuous Global Positioning System (CGPS) stations to determine the vertical movement of the land with respect to the International Terrestrial Reference Frame.

Observations collected by the sea level monitoring network are routinely processed into a range of quality-controlled data products. The monthly data report is the primary source of up-to-date information relating to these data products.






Figure 1. Pacific Sea Level Monitoring Network of SEAFRAME stations

Sea Level and Climate

Astronomical tides and weather conditions are largely responsible for daily perturbations in sea level, but over monthly, seasonal and longer timescales sea levels in the tropical Pacific are largely influenced by fluctuations in climate and ocean heat content across the Pacific.

The El Niño – Southern Oscillation climate cycle plays a key role in sea level variability. During El Niño sea levels are generally lower than normal across the western equatorial Pacific, as measured by the project's sea level network, in response to weaker than normal easterly Trade Winds, cooler than normal ocean temperatures and higher than normal barometric pressures in this region. On the other hand during La Niña the easterly Trade Winds are typically stronger than normal, ocean temperatures are warmer than normal and barometric pressures are lower than normal across the western Pacific, which often results in higher than normal sea levels at many of the project stations.

The sea level stations at PNG, Solomon Islands, Tuvalu and Samoa lie along a zone of convergent winds, known as the South Pacific Convergence Zone. Sea levels at these stations may become higher or lower than normal depending on the strength of these convergent winds or the shifting position of the convergence zone relative to its climatological mean. The sea level stations at Nauru and Kiribati lie very close to the equator and can both be influenced by sea level signals propagating along the equatorial waveguide.

A summary of recent and past climate conditions across the equatorial Pacific is provided by the Bureau of Meteorology in its monitoring of the El Niño – Southern Oscillation cycle at <http://www.bom.gov.au/climate/enso/>

Further climate information for Pacific Island countries is provided by the Climate and Ocean Monitoring and Prediction (COMP) Project under the Climate and Oceans Support Program in the Pacific (COSPPac).



October SEAFRAME Data

Monthly Sea Level and Environmental Data

The observed sea levels (Figure 3) are dominated by the daily oscillations of the tide. In most cases, the tide rises and falls twice per day (semi-diurnal), but at PNG and the Solomon Islands the tide tends to have a single high and low per day (diurnal). Where the tides follow a semi-diurnal pattern the greatest tidal variations are called spring tides, which tend to occur around the time of the full and new moons. There was a new moon on the 15th of October and a full moon on the 29th of October.

Gaps in the data are the result of instrumental errors or data retrieval problems and are discussed under Instrument Performance.

The residuals (Figure 4) are the differences between the observed sea levels and the astronomical tidal predictions. They highlight non-tidal sea level fluctuations, such as those due to the effects of weather or tsunamis. Tropical cyclones can produce storm surges where the combination of low barometric pressure and strong winds raise sea levels well above the predicted astronomical tides for a period of a day or more.

The non-tidal sea level fluctuations can be amplified or sustained by the shape of the harbour in which the gauge is located. Some of the SEAFRAME stations are located in harbours that exhibit 'sloshing' under certain conditions (a phenomenon referred to as a seiche), such as at PNG when the wind suddenly changes strength or direction, at FSM during smaller neap tides and at Nauru during strong westerly winds.

No significant storm surge was observed across the network during October. Sea levels remained, for the most part, close to predicted astronomical tides during October. At Nauru sea levels exceeded the predicted tide by 0.3m on 7th October 2012, but the tidal range at the time was quite small with the moon and sun being in quadrature. Small tsunami signals on the order of 10cm were observed at Samoa and Cook Islands following a magnitude Mw7.7 earthquake that struck at 03:04 UTC on 28th October 2012 in the northeast Pacific off the coast of British Columbia, Canada in the vicinity of Haida Gwaii (Queen Charlotte Islands).

The sea level residuals at all stations, to some degree, exhibit semi-diurnal or diurnal fluctuations, which last a few days or weeks and then disappear. If these fluctuations were to persist they would form part of the astronomical tide prediction and thus not appear as residuals. Consequently semi-diurnal and diurnal residual fluctuations will always be transient in nature.

The barometrically corrected residuals (Figure 5) have had the effect of atmospheric pressure fluctuations removed from the sea level residuals of Figure 4. The rule of thumb for the 'inverse barometer effect' is that a 1-hPa fall in the barometer, if sustained over a day or more, produces a 1-cm rise in the local sea level (within the area beneath the low pressure system).

The winds, temperatures and barometric pressures are plotted in Figure 6 to Figure 11. The incident

winds in Figure 7 follow the meteorological convention, that is, they point in the direction the wind is coming from. For example, the winds at Marshall Islands prevailed from the northeast for most of the month.

Air and water temperatures (Figure 9 and Figure 10) are plotted using the same vertical scale for the purpose of comparison. The air temperatures are seen to fluctuate over a much wider range than the water temperatures. At some sites (e.g. Marshall Islands) the water temperature shows almost no variation, although the air temperature varies by several degrees between night and day. At Nauru a twice-daily fluctuation in water temperature is related to interactions between tides and terrestrial (land-based) water discharging into the wharf area. The water temperature fluctuations there are usually more pronounced during the larger spring tides.

Barometric pressures (Figure 11) tend to fluctuate by around 3 hPa twice-daily at all stations as a result of atmospheric tides, which are largest in the tropical regions and reduce to near zero toward the

poles. The longer-term barometric pressure fluctuations that occur over periods of days to weeks are due to passing weather systems. These fluctuations tend to be larger at sites farther away from the equator such as Cook Islands and Tonga.

The meteorological data are put into perspective by Figure 12. In this figure, if an open circle falls above (below) a solid dot, a new maximum (minimum) for the particular month has been set. The data sets only include Pacific Sea Level Monitoring Project data, which have been collected since October 1992 when the first station was installed at Fiji. The data from FSM has only been collected since December 2001. Notable extremes recorded this month include record-high October air temperatures at Solomon Islands (32.3 °C) and Vanuatu (32.7 °C), while the monthly maximum air temperature at Kiribati and monthly maximum water temperature at Fiji equalled previous October records. Further sea level and meteorological statistical information is available at

<http://www.bom.gov.au/oceanography/projects/spslcm/data/monthly.shtml>



Mean Sea Level and Anomalies

Figure 13 shows the monthly mean sea levels, which are simple arithmetic averages of the sea levels, relative to an arbitrary zero. The figure shows that Tuvalu, for example, normally experiences an annual cycle of about 0.2 metres, reaching a peak around February or March. One effect of the El Niño of 1997-1998 was very low sea levels which disrupted the annual sea level cycle at many of the SEAFRAME stations.

Figure 14 shows the monthly mean sea level anomalies, which are the residuals after tides, annual and semi-annual seasonal cycles and linear slope have been removed by way of a harmonic tidal analysis of the complete record. The annual

cycle at Tuvalu (which has the largest consistent annual cycle) is quite notable in Figure 13 but less apparent in Figure 14. By removing the seasonal cycles, the anomalies help to bring out irregular features, such as lower than normal sea levels across the region during the 1997/98 El Niño.

The sea level anomalies for October 2012 reveal monthly mean sea levels were about 5cm higher than normal at Kiribati, Nauru and Fiji and 5cm lower than normal at Solomon Islands, Tuvalu and Vanuatu. Monthly sea levels at Marshall Islands, PNG, Samoa, Tonga and Cook Islands were near normal for this time of the year.

Overall Rate of Movement in Sea Level

Table 1 shows the overall rate of movement in sea level at individual Pacific stations based on the data so far collected at those sites. For many of the sites, the underlying data sets are around twenty years in length.

The overall rates of movement are updated every month by calculating the linear slope during the tidal analysis of all the data available at individual stations.

Please exercise caution in interpreting the overall rates of movement of sea level – the records are too short to be inferring long-term trends.

Table 1. Updated overall rates of sea level movement based on SEAFRAME data from installation through October, 2012.

Location	Latitude	Longitude	Date of first data	Rate (mm/yr)	Change in rate from previous month (mm/yr)
Cook Is	21°12'17.1"S	159°47'5.2"W	Feb 1993	+5.8	0.0
Tonga	21°8'12.5"S	175°10'50.5"W	Jan 1993	+9.0	0.0
Fiji	17°36'17.7"S	177°26'17.7"E	Oct 1992	+6.0	0.0
Vanuatu	17°45'19.2"S	168°18'27.7"E	Jan 1993	+4.9	-0.1
Samoa	13°49'36.4"S	171°45'40.7"W	Feb 1993	+7.6	+0.1
Tuvalu	8°30'8.9"S	179°11'42.6"E	Mar 1993	+3.9	-0.1
Kiribati	1°21'54.2"N	172°55'58.8"E	Dec 1992	+3.3	+0.1
Nauru	0°31'45.9"S	166°54'36.2"E	Jul 1993	+4.3	+0.1
Solomon Is.	9°25'44.1"S	159°57'19.3"E	Jul 1994	+7.9	-0.1
PNG	2°2'31.5"S	147°22'25.6"E	Sep 1994	+8.3	0.0
FSM	6°58'49.9"N	158°12'0.8"E	Dec 2001	+17.9	0.0
Marshall Is.	7°6'21.7"N	171°22'22.1"E	May 1993	+5.9	0.0



Barometric Pressure, Water Temperature and Air Temperature Anomalies

The anomalies of barometric pressure, water and air temperature are determined in the same manner as the sea level anomalies, except the linear slope is not calculated.

The barometric pressure anomalies (Figure 15) show substantially higher than normal barometric pressures were observed at SEAFRAME stations during the 1997-1998 El Niño. In comparison monthly average barometric pressures during October 2012 were much closer to normal.

Water temperature anomalies (Figure 16) show monthly water temperatures for October 2012 remained +1.0 °C warmer than normal at Nauru and slightly warmer than normal at Solomon Islands, Kiribati and Fiji. Slightly cooler than normal water temperatures were observed at PNG. Water temperatures at Marshall Islands, Samoa and Vanuatu through October were close to what is expected at this time of the year. Water

temperature data were unable to be collected at FSM, Tuvalu, Tonga and Cook Islands during October due to technical problems.

Air temperature anomalies (Figure 17) show warmer than normal conditions were observed at Kiribati and Samoa during October, where anomalies remained close to +1.0 °C. Warming conditions were also observed at Vanuatu, Tonga and Cook Islands during October with the development of air temperature anomalies exceeding +0.5 °C. Air temperatures at PNG, Solomon Islands, Tuvalu and Fiji were near normal for this time of the year while elsewhere (Marshall Islands, FSM, Nauru and Fiji) no air temperature data were able to be collected during October due to technical problems. Over the duration of the record the air temperature anomalies generally (although not always) follow the water temperature anomalies, which is an indication of the large influence the ocean has upon the climate of the Pacific Islands.

Instrument Performance

In Figure 18, which shows sea level data return, colour is used to distinguish five-year project phases. The number of missing days is noted in gaps in the bars.

Sea level data return from the network was only 85.5% during October 2012 (Table 2), due to significant station outages at Cook Islands and FSM. The station at Cook Islands was upgraded under the Observation Network Upgrade Project and returned to service on 20th October 2012, while

FSM is scheduled for a similar upgrade in February 2013.

Additional problems encountered with the ancillary sensors during October included the air temperature sensors at Marshall Islands, Nauru and Fiji, the water temperature sensors at Tuvalu and Tonga and the wind monitor at Tuvalu. The erroneous data received from these problematic sensors were removed from the quality-controlled archived records.

Table 2. Rates of sea level data return

Location	Installation Date	Data Return Since Installation (%)	Data Return in October 2012 (%)
Cook Is	Feb 1993	96.7	38.3
Tonga	Jan 1993	98.4	100
Fiji	Oct 1992	98.8	100
Vanuatu	Jan 1993	94.9	100
Samoa	Feb 1993	96.3	100
Tuvalu	Mar 1993	96.3	90.8
Kiribati	Dec 1992	93.8	100
Nauru	Jul 1993	92.6	96.6
Solomon Is.	Jul 1994	98.5	100
PNG	Sep 1994	93.2	100
FSM	Dec 2001	97.6	0.0
Marshall Is.	May 1993	98.1	100
Network Average		96.3	85.5



SEAFRAME Stations

SEAFRAME stations employ either a SUTRON or TELMET (for upgraded stations) programmable data logger, water level gauges and other sensors. The data logger and associated electronics are normally housed in fibreglass huts. A sketch of a typical SEAFRAME station is shown in Figure 2.

Water level sensors include:

1. Primary water level using a Bartex 'AQUATRAK' acoustic-in-air sensor,
2. Secondary water level (or backup) using a Druck pressure transducer mounted close to the seabed, and

3. Tertiary water level using a Vega-puls radar sensor mounted above the water (at upgraded sites).

For SUTRON stations, the water level samples are averaged over three minutes and logged every six minutes, while meteorological sensors are logged on an hourly basis. With the upgraded TELMET stations, the water level samples are averaged over one minute and, together with meteorological data, logged every minute. Appropriate weighted-average and time-centred data is computed remotely which conforms to the SUTRON algorithm. Both SUTRON and TELMET data loggers have the memory capacity to store approximately one month of data




Figure 2. Schematic diagram of a SEAFRAME sea level monitoring station.

Observation Network Upgrade Project

The Observation Network Upgrade Project (ONUP) is scheduled to upgrade all Pacific SEAFRAME stations by Dec-2013 with modernised TELMET

data loggers, real-time satellite communications and additional radar-type water level sensors. The status of the station upgrades is given in Table 3.

Table 3. Schedule of SEAFRAME station equipment upgrades.

Location	Latitude	Longitude	SUTRON Installation Date	TELMET Upgrade Date
Cook Is	21°12'17.1"S	159°47'5.2"W	Feb 1993	Oct 2012
Tonga	21°8'12.5"S	175°10'50.5"W	Jan 1993	Mar 2011
Fiji	17°36'17.7"S	177°26'17.7"E	Oct 1992	Jun 2011
Vanuatu	17°45'19.2"S	168°18'27.7"E	Jan 1993	May 2012
Samoa	13°49'36.4"S	171°45'40.7"W	Feb 1993	Aug 2011
Tuvalu	8°30'8.9"S	179°11'42.6"E	Mar 1993	To be upgraded
Kiribati	1°21'54.2"N	172°55'58.8"E	Dec 1992	Oct 2011
Nauru	0°31'45.9"S	166°54'36.2"E	Jul 1993	To be upgraded
Solomon Is.	9°25'44.1"S	159°57'19.3"E	Jul 1994	Nov 2011
PNG	2°2'31.5"S	147°22'25.6"E	Sep 1994	Aug 2012
FSM	6°58'49.9"N	158°12'0.8"E	Dec 2001	To be upgraded
Marshall Is.	7°6'21.7"N	171°22'22.1"E	May 1993	To be upgraded



Further Information

Online Resources

COSPPac Web site: <http://www.bom.gov.au/cosppac/>

PSLMP Web site: <http://www.bom.gov.au/oceanography/projects/spslcmp/spslcmp.shtml>

ENSO Wrap-Up - El Niño / La Niña information: <http://www.bom.gov.au/climate/enso/>

Geoscience Australia South Pacific Regional GNSS Network (Levelling Survey and Continuous GPS Monitoring):
<http://www.ga.gov.au/earth-monitoring/geodesy/gnss-networks.html>

Acknowledgement

The Monthly Data Report is prepared for AusAID by the National Tidal Centre (NTC), Bureau of Meteorology under the Pacific Sea Level Monitoring (PSLM) Project, Climate and Oceans Support Program in the Pacific (COSPPac).

Further enquiries about the Monthly Data Report may be made to NTC at:

National Tidal Centre
Bureau of Meteorology
PO Box 421
Kent Town SA 5067
Tel: (+618) (08) 8366 2730
Email: ntc@bom.gov.au
Website: <http://www.bom.gov.au/oceanography/tides.shtml>

Please also note the following:

While care has been taken in the collection, analysis, and compilation of the data, it is supplied on the condition that neither the Commonwealth of Australia nor NTC shall be liable for any loss or injury whatsoever arising from the use of the data. Copyright for material contained in this document is held by the Commonwealth of Australia.

Appendix 1: SEAFRAME Data Figures



SIX MINUTE WATER LEVEL OBSERVATIONS (m)

October 2012




Figure 3. Sea level observations during October 2012.

SIX MINUTE RESIDUAL WATER LEVELS (m)

October 2012




Figure 4. Residual sea levels during October 2012.



SIX MINUTE RESIDUALS ADJUSTED FOR ATMOSPHERIC PRESSURE (m)

October 2012




Figure 5. Residual sea levels adjusted for atmospheric pressure during October 2012.

HOURLY WIND SPEEDS (m/s)

October 2012




Figure 6. Wind speeds during October 2012.



HOURLY INCIDENT WINDS (m/s, deg True)




Figure 7. Incident winds during October 2012.

HOURLY MAXIMUM WIND GUSTS (m/s)

October 2012




Figure 8. Wind gusts during October 2012.



HOURLY AIR TEMPERATURES (°C)

October 2012




Figure 9. Air temperatures during October 2012.

HOURLY WATER TEMPERATURES (°C)

October 2012




Figure 10. Water temperatures during October 2012.



HOURLY ATMOSPHERIC PRESSURE (hPa)

October 2012







Figure 11. Atmospheric pressures during October 2012.

Comparison of October 2012 Max, Min & Mean with Long Term October Values



Stations

- 1 - Marshall Islands
- 2 - Federated States of Micronesia
- 3 - Papua New Guinea
- 4 - Solomon Islands
- 5 - Kiribati
- 6 - Nauru
- 7 - Tuvalu
- 8 - Samoa
- 9 - Vanuatu
- 10 - Fiji
- 11 - Tonga
- 12 - Cook Islands

- October 2012 Maximum
- October 2012 Mean
- October 2012 Minimum

- Long Term October Maximum
- Long Term October Mean
- Long Term October Minimum

Figure 12. Comparison of October 2012 ancillary data with long term October values



MONTHLY MEAN SEA LEVELS TO OCTOBER 2012 (m)

The zero line represents an arbitrary fixed offset from the zero of the tide gauge.




Figure 13. Monthly mean sea levels to October 2012

SEA LEVEL ANOMALIES THROUGH OCTOBER 2012 (m)




Figure 14. Monthly sea level anomalies to October 2012.



BAROMETRIC PRESSURE ANOMALIES THROUGH OCTOBER 2012 (hPa)




Figure 15. Monthly barometric pressure anomalies to October 2012.

WATER TEMPERATURE ANOMALIES THROUGH OCTOBER 2012 (°C)




Figure 16. Monthly water temperature anomalies to October 2012.



AIR TEMPERATURE ANOMALIES THROUGH OCTOBER 2012 (°C)




Figure 17. Monthly air temperature anomalies to October 2012.

SEA LEVEL DATA RETURN

THE NUMBER OF DAYS OF GAP ARE INDICATED
GAPS INCLUDE TRANSMISSION, POWER AND LOGGER FAILURE




Figure 18. Sea level data return.